

Exact Solution to the Moment Problem for the XY Chain

N.S. Witte*

*Research Centre for High Energy Physics,
School of Physics, University of Melbourne,
Parkville, Victoria 3052, AUSTRALIA.*

February 7, 2008

Abstract

We present the exact solution to the moment problem for the spin-1/2 isotropic antiferromagnetic XY chain with explicit forms for the moments with respect to the Néel state, the cumulant generating function, and the Resolvent Operator. We verify the correctness of the Horn-Weinstein Theorems, but the analytic structure of the generating function $\langle e^{-tH} \rangle$ in the complex t -plane is quite different from that assumed by the t -Expansion and the Connected Moments Expansion due to the vanishing gap. This function has a finite radius of convergence about $t = 0$, and for large t has a leading descending algebraic series $E(t) - E_0 \sim At^{-2}$. The Resolvent has a branch cut and essential singularity near the ground state energy of the form $G(s)/s \sim B|s+1|^{-3/4} \exp(C|s+1|^{1/2})$. Consequently extrapolation strategies based on these assumptions are flawed and in practise we find that the CMX methods are pathological and cannot be applied, while numerical evidence for two of the t -expansion methods indicates a clear asymptotic convergence behaviour with truncation order.

PACS: 05.30.-d, 11.15.Tk, 71.10.Pm, 75.10.Jm

*E-mail: nswphysics.unimelb.edu.au

1 Motivation

Some novel non-perturbative formalisms based on a moment approach have been proposed recently, the t -Expansion[1] and the Connected Moments Expansion (CMX)[2], and applied in a number of areas ranging from quantum chemistry[2, 3, 4, 5, 6], lattice gauge theories [7, 8, 9, 10, 11, 12, 13] to condensed matter problems[14, 15, 16, 17, 18]. These formalisms take their starting point from a finite sequence of low order connected Hamiltonian moments, or cumulants $\{\nu_n : n = 1 \dots\}$, and appeal to ostensibly exact theorems for ground state properties in order to extract appropriate averages. In all these applications, the sequence of cumulants was finite and relatively small, and the effects of truncation and the rate of convergence was completely unknown. This issue has particular practical importance because in many of the applications the convergence and final estimates were rather poor considering the effort expended in calculating a large number of moments. As rough rule of thumb, the initial and low order estimates were very good and the situation looked promising but when taken to higher order there was not a significant improvement, and in some cases the convergent behaviour seemed to be arrested[17, 19, 18]. The expectation of an accurate, perhaps arbitrarily and controllably accurate, method has to date seemed to be unfounded. The source of these problems had not been properly identified nor any remedy proposed. Here we give the first example of an exact solution to a moment problem which can answer some of these questions.

Furthermore it was not known whether these techniques were capable of yielding exact results in any case. This is a natural expectation since any formalism, applicable to general problems, must have a set of simple, but nontrivial solvable cases. There was another model[20], of quadratically coupled harmonic oscillators, for which the exact ground state wavefunction and density matrices are known but the moment problem was not exactly solved in that a general cumulant was not given. We present the solution to the moment problem for the isotropic, one-dimensional antiferromagnetic spin $\frac{1}{2}$ XY Model. Our solution also demonstrates an alternative notion of solvability, from those concepts and methods usually associated with the known solvable models.

In the t -Expansion, one constructs the cumulant generating function from its Taylor series expansion about $t = 0$,

$$\langle e^{tH} \rangle = \exp \left\{ \sum_{n=1}^{\infty} \nu_n \frac{t^n}{n!} \right\} \equiv e^{F(t)} , \quad (1)$$

where all moments are calculated with respect to some trial state, with the proviso that the overlap with exact ground state is nonzero. Then the Horn-Weinstein Theorem[1] states that the ground state energy and first excited state gap (within the same sector as the ground state) are given by the following limits

$$\begin{aligned} E_0 &= \lim_{t \rightarrow \infty} E(t) \\ E_1 - E_0 &= \lim_{t \rightarrow \infty} -\frac{d^2 E(t)/dt^2}{dE(t)/dt} , \end{aligned} \quad (2)$$

where $E(t) \equiv F'(-t)$. Other ground state averages can be found as well by computing the appropriate generalised connected moments[1] and performing the same extrapolation. This theorem was “proved” in Ref. ([1]) by assuming a simple discrete point

spectrum for the system, even in the thermodynamic limit, and these kinds of assumptions continue to be commonly used. The question immediately arises of how this extrapolation can be carried out and several methods[21] are employed. These are

Padé Analysis:

$E(t)$ is assumed to be representable by a $[L/M]$ Padé approximant with $L = M$ i.e. it has only separable poles and zeros of finite order in any finite region of the complex t -plane, and the extrapolation $t \rightarrow \infty$ made directly,

D-Padé Analysis:

$dE(t)/dt$ is assumed to be representable by a $[L/M]$ Padé approximant with $M \geq L + 2$, on the basis of the same considerations as above, and the ground state energy is found by numerically integrating this

$$E(\infty) - E(0) = \int_0^\infty ds E'(s) , \quad (3)$$

Laplace Method/Resolvent Analysis:

The Resolvent

$$G(u)/u \equiv \langle \frac{1}{1 + uH} \rangle , \quad (4)$$

is assumed to have a pole structure of the form

$$G(u)/u \sim \frac{b_0}{1 + uE_0} + \frac{b_1}{1 + uE_1} + \dots , \quad (5)$$

so that a $[L/M]$ Padé analysis with $M = L + 1$ can be made and the pole found,

Inversion Analysis:

One inverts the $E(t)$ function, forms $dt(E)/dE$ and makes a Padé analysis of this. The ground state energy is the location of a pole in this derivative, from

$$\frac{dt}{dE} \sim \frac{C}{E - E_0} , \quad (6)$$

“ E of F ”/Partition Function:

The function $\delta(t) = 1 - \langle e^{-tH} \rangle$ is taken as the independent variable instead of t . One inverts the $\delta(t)$ function and makes a diagonal Padé analysis of $E(\delta)$ and extracts the ground state energy from

$$E_0 = \lim_{\delta \rightarrow 1^-} \frac{1}{(1 - \delta) \frac{dt}{d\delta}} . \quad (7)$$

The Connected Moments Expansion, of which there are many variants, is derived by assuming

$$E(t) \sim E_0 + \sum_{j=1}^n A_j e^{-b_j t} . \quad (8)$$

The variants give the following explicit forms for the ground state energy

CMX-HW:

The first method[2] gives the ground state energy up the n th order as

$$E_0 = \nu_1 - \sum_{i=1}^n \frac{S_{i,2}^2}{\prod_{j=1}^i S_{j,3}} , \quad (9)$$

where the S -variables are recursively generated by

$$\begin{aligned} S_{1,k} &= \nu_k \\ S_{i+1,k} &= S_{i,k} S_{i,k+2} - S_{i,k+1}^2 , \end{aligned} \quad (10)$$

CMX-SD:

Another method[20] gives the ground state energy up the n th order as

$$E_0 = \omega_{1,n} , \quad (11)$$

where the ω -variables are recursively generated by

$$\begin{aligned} \omega_{j,1} &= \nu_j \\ \omega_{j,k+1} &= \omega_{j,k} - 2 \frac{\omega_{j+1,k} \omega_{2,k}}{\omega_{3,k}} + \left(\frac{\omega_{2,k}}{\omega_{3,k}} \right)^2 \omega_{j+2,k} , \end{aligned} \quad (12)$$

CMX-LT:

This method described in Ref. [20, 22] gives

$$E_0 = \nu_1 - (\nu_2, \nu_3, \dots, \nu_n) \begin{pmatrix} \nu_3 & \nu_4 & \cdots & \nu_{n+1} \\ \nu_4 & \nu_5 & \cdots & \nu_{n+2} \\ \vdots & \vdots & \ddots & \vdots \\ \nu_{n+1} & \nu_{n+2} & \cdots & \nu_{2n-1} \end{pmatrix}^{-1} \begin{pmatrix} \nu_2 \\ \nu_3 \\ \vdots \\ \nu_n \end{pmatrix} . \quad (13)$$

There is also another variant, called the AMX[23], which is very similar to the CMX-HW method. The CMX, in addition to its convergence difficulties, often presents another, although less serious problem. Usually the cumulants are nonlinear functions of some coupling constant, and the resulting ground state energy estimate above truncated at some order above would be rational function of the constant. In these problems[24, 25] an unphysical pole would appear for real coupling values. This pole was found to be exactly cancelled[23] if one included the next order, but a next order is not always available.

2 The Model

We follow the conventions and notations of Ref. [26] as closely as possible in what follows. For the N site chain, with N even but not a multiple of 4, the Hamiltonian is taken to be

$$H = \sum_{i=1}^N [S_i^x S_{i+1}^x + S_i^y S_{i+1}^y] , \quad (14)$$

where the labels $i, j, \dots$ index the sites. Following the standard treatments a transformation from spin operators to a mixed Fermi-Bose operator algebra is performed

$$a_i^\dagger, a_i \equiv S_i^x \pm iS_i^y, \quad (15)$$

and then a Wigner-Jordan transformation to a pure Fermionic representation

$$\begin{aligned} c_i &\equiv e^{\pi i \sum_{j=1}^{i-1} a_j^\dagger a_j} a_i \\ c_i^\dagger &\equiv a_i^\dagger e^{-\pi i \sum_{j=1}^{i-1} a_j^\dagger a_j}, \end{aligned} \quad (16)$$

so that the Hamiltonian is now

$$H = \frac{1}{2} \sum_{i=1}^N \left[c_i^\dagger c_{i+1} + c_{i+1}^\dagger c_i \right]. \quad (17)$$

In doing so we have neglected the boundary terms arising from the exact mapping of the spin or a-cyclic problem, and will only consider the c-cyclic problem $c_{N+1} = c_1$. We take this course because we don't want to complicate matters unnecessarily and are only interested in the thermodynamic limits.

Proceeding with the diagonalisation of this quadratic form via the canonical transformation

$$\begin{aligned} \eta_p &= \sum_i \left[\frac{\phi_{pi} + \psi_{pi}}{2} c_i + \frac{\phi_{pi} - \psi_{pi}}{2} c_i^\dagger \right] \\ \eta_p^\dagger &= \sum_i \left[\frac{\phi_{pi} + \psi_{pi}}{2} c_i^\dagger + \frac{\phi_{pi} - \psi_{pi}}{2} c_i \right], \end{aligned} \quad (18)$$

with transformation coefficients

$$\begin{aligned} \phi_{pi} &= \sqrt{\frac{2}{N}} \sin pi & p > 0 \\ &= \sqrt{\frac{2}{N}} \cos pi & p \leq 0 \\ \psi_{pi} &= \frac{\cos p}{\Lambda_p} \phi_{pi}, \end{aligned} \quad (19)$$

where the quasiparticle energy $\Lambda_p = |\cos p|$ and the momentum values, labelled by $p, q, \dots$ are

$$p = \frac{2\pi m}{N} \quad m = -N/2, \dots, 0, 1, \dots, N/2 - 1. \quad (20)$$

The Hamiltonian then becomes

$$H = \sum_p \Lambda_p \eta_p^\dagger \eta_p - \frac{1}{2} \sum_p \Lambda_p. \quad (21)$$

The transformation coefficients are orthonormal in the sense

$$\begin{aligned} \phi_p \cdot \phi_{p'} &= \delta_{pp'} \\ \psi_p \cdot \psi_{p'} &= \delta_{pp'} \\ \phi_p \cdot \psi_{p'} &= 0. \end{aligned} \quad (22)$$

In all the moment formalisms a choice has to be made as what trial state the moments of the Hamiltonian are computed with respect to and it is this choice which is the crucial physical one. The only general requirement is that, for ground state properties at least, the trial state should have a non-zero overlap with the true ground state. In all the previous and approximate treatments of the the XY chain, and most of the other antiferromagnetic spin models, the trial state was taken to be the classical Néel state. This state is in fact, as we shall quantify, as far from the true ground state as one could possibly be but its simplicity allows for ease of moment computation, certainly compared to alternatives. So as a “natural” choice we proceed by only considering this choice.

In the spin picture the Néel state $|N\rangle$ will be represented by

$$|\uparrow\downarrow\uparrow\downarrow\ldots\uparrow\downarrow\uparrow\downarrow\rangle, \quad (23)$$

and in the fermionic c -picture by

$$|1010\ldots1010\rangle. \quad (24)$$

The defining equations for the Néel state are

$$\left. \begin{aligned} c_{2i+1}^\dagger |N\rangle &= 0 \\ c_{2i} |N\rangle &= 0 \end{aligned} \right\} \quad \forall i \quad (25)$$

and using the transformation coefficients, Eq. (19), and the spectral points, Eq.(20), these can be shown to be equivalent to the following equations in terms of the quasi-particle operators

$$\left. \begin{aligned} (\eta_p - \eta_{\bar{p}}^\dagger) |N\rangle &= 0 \\ (\eta_p^\dagger + \eta_{\bar{p}}) |N\rangle &= 0 \end{aligned} \right\} \quad \forall p \in Z. \quad (26)$$

Here the spectrum of allowed momentum values have been divided up into the set

$$Z \equiv [-\pi, -\pi/2) \cup (0, \pi/2), \quad (27)$$

and its complement $\bar{Z}$, and there is a bijection from Z to $\bar{Z}$ defined by

$$\begin{aligned} \bar{q} &\equiv -q - \pi & q &\leq 0 \\ &\equiv -q + \pi & q &> 0. \end{aligned} \quad (28)$$

If we denote the vacuum state for the quasiparticles by $|0\rangle$ then the Néel state can be expressed in terms of the quasiparticles by

$$|N\rangle = 2^{-N/4} \left\{ 1 + \sum_{q \in Z} \eta_q^\dagger \eta_{\bar{q}}^\dagger + \sum_{q_1 < q_2 \in Z} \eta_{q_1}^\dagger \eta_{\bar{q}_1}^\dagger \eta_{q_2}^\dagger \eta_{\bar{q}_2}^\dagger \right. \\ \left. \ldots + \eta_{q_1}^\dagger \eta_{\bar{q}_1}^\dagger \eta_{q_2}^\dagger \eta_{\bar{q}_2}^\dagger \ldots \eta_{q_{N/2}}^\dagger \eta_{\bar{q}_{N/2}}^\dagger \right\} |0\rangle, \quad (29)$$

and because of the anticommutation relations for the quasiparticle operators

$$\begin{aligned} |N\rangle &= 2^{-N/4} \exp \left(\sum_{q \in Z} \eta_q^\dagger \eta_q^\dagger \right) |0\rangle \\ &= \prod_{q \in Z} \frac{1 + \eta_q^\dagger \eta_q^\dagger}{\sqrt{2}} |0\rangle . \end{aligned} \quad (30)$$

One can easily show that this state satisfies the defining Eq. (26). Basically the Néel state is created from the quasiparticle vacuum state by pairs of excitations whose momenta sum to $\pm\pi$.

We will have need of some spectral sums defined on Z and these are

$$\begin{aligned} \sum_{q \in Z} \Lambda_q^{2n} &= \frac{(2n)!}{2^{2n+1}(n!)^2} N \\ \sum_{q \in Z} \Lambda_q &= \frac{1}{\sin(\pi/N)} . \end{aligned} \quad (31)$$

3 Generating Function

We now show how to find the exact form for the generating function

$$\langle e^{tH} \rangle \equiv \langle N | e^{tH} | N \rangle = \sum_{n=0}^{\infty} \frac{\mu_n t^n}{n!} , \quad (32)$$

where the moments $\mu_n \equiv \langle H^n \rangle$. We will establish this result for a finite system of size N although ultimately the thermodynamic limit will be taken. It is straightforward to show a direct expression for a general moment

$$\begin{aligned} \mu_n &= 2^{-N/2} \left\{ E_0^n + \sum_{q \in Z} (E_0 + 2\Lambda_q)^n + \frac{1}{2!} \sum_{q_1 \neq q_2 \in Z} (E_0 + 2\Lambda_{q_1} + 2\Lambda_{q_2})^n \right. \\ &\quad \left. \dots + (E_0 + 2\Lambda_{q_1} + \dots + 2\Lambda_{q_{N/2}})^n \right\} . \end{aligned} \quad (33)$$

in its symmetrical form. Then our initial form for the generating function is

$$\begin{aligned} \langle e^{tH} \rangle &= 2^{-N/2} e^{E_0 t} \left\{ 1 + \sum_{q \in Z} e^{2\Lambda_q t} + \frac{1}{2!} \sum_{q_1 \neq q_2 \in Z} e^{2t(\Lambda_{q_1} + \Lambda_{q_2})} \right. \\ &\quad \left. \dots + e^{2t(\Lambda_{q_1} + \dots + \Lambda_{q_{N/2}})} \right\} . \end{aligned} \quad (34)$$

For simplicity let us define the following sum

$$I_n \equiv \sum_{q \in Z} e^{2nt\Lambda_q} , \quad (35)$$

and denote the general p -th term of the Eq. (34) by $g_p/p!$ after expanding it. This would now be

$$\frac{g_p}{p!} = \frac{1}{p!} \sum_{q_1 \neq \dots \neq q_p \in Z} e^{2t\Lambda_{q_1}} \dots e^{2t\Lambda_{q_p}} . \quad (36)$$

Such sums over mutually distinct spectral points are recognisable as the cycle index of the Symmetric Group S_p . Consequently we have the partition sum formula

$$\frac{g_p}{p!} = \sum_{\sum i_i = p} \frac{1}{i_1! 1^{i_1} i_2! 2^{i_2} \dots i_p! p^{i_p}} (-)^{p+\sum i_i} I_1^{i_1} \dots I_p^{i_p} , \quad (37)$$

and find immediately the exponential sum, constituting the generating function, is

$$\sum_{p=0}^{\infty} \frac{g_p}{p!} = \exp \left\{ \sum_{n=1}^{\infty} \frac{(-)^{n+1}}{n} I_n \right\} . \quad (38)$$

Each term $g_p/p!$ is an elementary symmetric polynomial in the variables $z_p \equiv e^{2t\Lambda_p}$ and Eq. (37) is just the way of re-expressing this in terms of the power sums $I_n = \sum_p z_p^n$.

The generating function is then expressible in the following sequence of closed forms

$$\begin{aligned} \langle e^{tH} \rangle &= 2^{-N/2} e^{tE_0} \exp \left\{ \frac{1}{2} \log \prod_q (1 + e^{2t\Lambda_q}) \right\} \\ &= \prod_{q \in Z} \cosh(t\Lambda_q) \\ &= \prod_{k=0}^{\infty} \prod_{q \in Z} \left(1 + \frac{1}{\pi^2(k+1/2)^2} t^2 \Lambda_q^2 \right) \\ &\xrightarrow{N \rightarrow \infty} \exp \left\{ \frac{N}{4\pi} \int_{-\pi}^{\pi} dq \log \cosh(t\Lambda_q) \right\} . \end{aligned} \quad (39)$$

This leads to the generating function $F(t)$ associated with the t -expansion

$$F(t) = \frac{N}{\pi} \int_0^{\pi/2} dq \log \cosh(t\Lambda_q) , \quad (40)$$

which is identical to the form of the free energy[26] for this system of noninteracting quasiparticles with zero chemical potential, with the identification $\frac{1}{2}\beta \rightarrow t$. The corresponding density $f(t)$, defined by $F(t) \equiv Nf(t)$ is,

$$e^{f(t)} = \prod_{k=0}^{\infty} \left[\frac{1}{2} + \frac{1}{2} \sqrt{1 + \frac{t^2}{\pi^2(k+1/2)^2}} \right] . \quad (41)$$

This function is also the cumulant generating function, so from the taylor series expansion about $t = 0$ we have

$$\begin{aligned} \nu_{2n+1} &= 0 & n &\geq 0 \\ \nu_{2n} &= \frac{2^{2n}-1}{4n} \frac{(2n)!}{(n!)^2} B_{2n} N & n &\geq 1 , \end{aligned} \quad (42)$$

where B_{2n} are the standard Bernoulli numbers. These cumulants coincide exactly with all the low order ones[27] computed directly by combinatorial enumeration on a computer.

There is also a simpler, more useful expression for the moments, other than that given by Eq. (33). Using the infinite product expansion for the generating function and expanding it in t^2 , one collects all factors contributing to t^{2n} term. By picking n such factors, all of which must be distinct,

$$\frac{\mu_{2n}}{(2n)!} = \frac{1}{n!} \left(\frac{4}{\pi^2} \right)^n \sum_{(k_1, q_1) \neq \dots \neq (k_n, q_n)} \frac{\Lambda_{q_1}^2}{(2k_1+1)^2} \cdots \frac{\Lambda_{q_n}^2}{(2k_n+1)^2} , \quad (43)$$

and using

$$\sum_{k=0}^{\infty} \frac{1}{(2k+1)^{2i}} \sum_{q \in \mathbb{Z}} \Lambda_q^{2i} = \frac{\pi^{2i}(2^{2i}-1)}{2^{2i+2}(i!)^2} |B_{2i}|N , \quad (44)$$

one can apply the cycle index formula to arrive at

$$\mu_{2n} = (2n)! \sum_{\sum l_i = n} \prod_{l=1}^n \frac{1}{i_l!} \left(\frac{(-)^{l+1}(2^{2l}-1)}{4l(l!)^2} |B_{2l}|N \right)^{i_l} . \quad (45)$$

4 Analyticity and Analysis

In this part we analyse our results for the generating function and its closely related functions. The cumulant generating function $F(t)$ and its derived functions have a finite radius of convergence about $t = 0$ of $|t| < \pi/2$, and which is evident from the infinite product form of Eq. (41) in having branch points at $t = \pi i(k + 1/2)$ for all integer k . This finite radius of convergence about $t = 0$ will then be a source of problems for extrapolation methods, even though the function itself is analytic in the neighbourhood of $\Re(t) \geq 0$.

Focussing on

$$E(t) = -\frac{N}{\pi} \int_0^{\pi/2} dq \Lambda_q \tanh(t\Lambda_q) , \quad (46)$$

we verify the Horn-Weinstein Theorem is exactly satisfied

$$\lim_{t \rightarrow \infty} E(t) = -\frac{N}{\pi} . \quad (47)$$

Now the first excited state has an energy $X_1 = E_0 + 2\Lambda_q$ for a value of

$$q = \pm \frac{\pi}{2} \left(1 \pm \frac{2}{N} \right) , \quad (48)$$

so that $X_1 - E_0 \sim 2\pi/N$ and that the gap vanishes like $O(1/N)$. The Horn-Weinstein theorem for the gap also gives zero. Finally the expression for the overlap[28]

$$|\langle N|0 \rangle|^2 = \exp \left\{ - \int_0^{\infty} dt [E(t) - E(\infty)] \right\} , \quad (49)$$

can also be shown to give the correct answer of $2^{-N/2}$. The vanishing of this in the thermodynamic limit is another signal of the problems with using the Néel state.

However while the specific predictions of the theorem are correct the basis for these are not, as will be evident in what follows. We require for this the large t development of the function $E(t)$ and this can be obtained in a number of ways. Using the Mellin Inversion Theorems one can get the leading series only, so we adopt a direct expansion of the integrand. This yields

$$E(t) - E_0 = \frac{N}{2\pi} \sum_{m=1}^{\infty} \frac{\Gamma(m-1/2)}{m!\Gamma(1/2)} (1-2^{1-2m}) |B_{2m}| \left(\frac{\pi}{t}\right)^{2m} \\ + \frac{2N}{\pi} \sum_{n=1}^{\infty} (-)^n e^{-2nt} \sum_{m=0}^{\infty} \frac{(1/2)_m (2m+1)!}{m!} \sum_{k=0}^{2m+1} \frac{1}{k!} (2nt)^{-2m-2+k} . \quad (50)$$

Several comments are in order here. The above m sums are separately divergent sums, and consequently these expansions are asymptotic, but when combined the result is convergent. The leading terms are algebraic, and the exponential terms are subdominant, with a numerical evaluation

$$E(t) - E_0 \sim \frac{N}{2\pi} \left\{ \frac{1}{12} \left(\frac{\pi}{t}\right)^2 + \frac{7}{960} \left(\frac{\pi}{t}\right)^4 + \dots \right\} , \quad (51)$$

which is at complete variance with common assumptions discussed above. This is basically due to the vanishing spectral gaps. The specific leading exponent of 2 and not 1 in the descending series of t is also significant and different from the Padé analysis assumptions. A diagonal Padé analysis of the Taylor series for $E(t)$ is not possible because the ν_{2n+1} are zero, and so the $t \rightarrow \infty$ limit either doesn't exist or is trivially zero.

However a D-Padé analysis is possible as $[M/M+2]$ approximants can be constructed for $E'(t)$. We display in Figure 1 the error for the ground state energy, computed using a series of D-Padé approximants for M values up to 24. It is clear that while there is initial convergence, this eventually ceases and no further improvement is evident. And the best result is never any better than 10^{-4} . Furthermore there are some gaps in our data where the Padé approximant had singularities close to the positive real axis which made the numerical integration algorithm unreliable. There are two sources of error in this analysis, the first being the error due to the finite Padé approximants of the integrand, and the second due to the numerical integration. We consider the latter error to be small (much smaller than 10^{-4}) and relatively uniform across the series so that most of the variation is due to the truncation.

The Inversion Method is also flawed here in that the leading order factor for $t(E)$ has a branch cut singularity, not a pole

$$t(E) \sim \left(\frac{\pi}{24}N\right)^{1/2} \frac{1}{\sqrt{E - E_0}} . \quad (52)$$

We have done an inversion analysis, using a full range of Padé approximants $[L/M]$ to Eq. (6), and present the error for the ground state energy in Figure 2 against L . There are some points not evident in this Figure, and that is the root of the denominator

polynomial closest to the exact ground state energy was not necessarily real, nor the one with the lowest real part. Here it is also clear that there is good initial convergence but the improvement ceases and in fact gets worse. There are some points much closer to the true answer than the rest but it is also true that there is clustering around this point and there is no way of uniquely distinguishing this point. Also going to higher order doesn't lead to any improvement.

The Resolvent can be exactly written down as

$$G(s)/s = \int_0^\infty dx \exp \left\{ \frac{N}{\pi} \left[-x + \int_0^{\pi/2} dq \log \cosh(sx\Lambda_q) \right] \right\} , \quad (53)$$

in terms of a scaled inverse energy $s = uN/\pi$, although not explicitly evaluated. Instead we present a saddle point approximation to this integral for $N \rightarrow \infty$ which yields

$$G(s)/s \sim \left\{ \frac{N}{2\pi^2} s^2 \int_0^{\pi/2} dq \Lambda_q^2 \text{sech}^2(sx_0\Lambda_q) \right\}^{-1/2} \times \exp \left\{ \frac{N}{\pi} \left[-x_0 + \int_0^{\pi/2} dq \log \cosh(sx_0\Lambda_q) \right] \right\} , \quad (54)$$

where the saddle point x_0 is given by the functional equation

$$1 = s \int_0^{\pi/2} dq \Lambda_q \tanh(sx_0\Lambda_q) . \quad (55)$$

The inverse energy lies in the range $|s| > 1$ and as $s \rightarrow -1$ then the point $x_0 \rightarrow \infty$. Using the asymptotic form for this integral, namely from the large t analysis Eq. (51), one finds in the neighbourhood of the ground state energy

$$x_0 \sim \frac{\pi}{2\sqrt{6}} |s + 1|^{-1/2} , \quad (56)$$

and that the Resolvent behaves like

$$G(s)/s \sim 2^{-N/2} \left(\frac{4.6^{3/2}}{\pi^{7/2}} N \right)^{-1/2} |s + 1|^{-3/4} \exp \left(\frac{N}{2\sqrt{6}} |s + 1|^{1/2} \right) . \quad (57)$$

So that we have a branch cut and essential singularity, and not simple separated poles.

For the “ E of F ” analysis the inversion of the function $\delta(t)$ for $\delta \rightarrow 1^-$ has the following leading order term

$$t = \frac{\pi}{2} \log 2 + \frac{\pi}{N} \log(1 - \delta) + O(1/\log(1 - \delta)) , \quad (58)$$

and consequently the energy function $E(\delta)$ has the leading term as $\delta \rightarrow 1^-$ (with fixed N) of

$$E(\delta) - E_0 \sim \frac{N^3}{24\pi} \frac{1}{[\log(1 - \delta)]^2} , \quad (59)$$

which is not a rational function of δ .

The Connected Moments Expansions actually exhibit pathological behaviour in this Model in that for both the CMX-HW and CMX-SD the recursion can't even get started because $\nu_3 = 0$. And for the CMX-LT when the dimension of the matrix to be inverted is odd then it is exactly singular, and when it is even the inner product is exactly zero, which is again due to ν_3 vanishing. This behaviour is typical in that the expansions work better when ν_3 is large, say relative to $\nu_2^{3/2}$.

5 Conclusions

We have given the exact solution to the moment problem for the one dimensional isotropic spin-half antiferromagnetic XY Model and explicitly found expressions for the moments, the generating function and the Resolvent. With this Model we have shown the Theorems used by the t -Expansion and Connected Moment Expansion are correct but that the assumptions upon which they are based, are not true. In fact we find leading order algebraic decay in $E(t) - E_0$ of second degree and not sums of exponential decay terms, that the inversion of $E(t)$ has a leading order branch cut, and that the Resolvent in the neighbourhood of the ground state energy has branch cut and essential singularities. For this Model we find the CMX methods and some of the t -expansion extrapolation methods to give pathological results, and the two t -expansion methods that can be applied give clear indications of having asymptotic convergence behaviour with respect to the truncation order. The analytic properties of this Model are at complete variance with commonly held assumptions made by the methods and shows why extrapolation and expansion procedures based on these assumptions have poor convergence. However we optimistically predict that revision of the extrapolation methods, along lines indicated by our results, should dramatically improve the convergence characteristics of the t -Expansion and CMX-type methods. Although the XY Model is perhaps the most difficult case to treat with moment methods employing the Néel state as the trial state, being the most extreme quantum limit of the anisotropic antiferromagnetic Heisenberg Models, a number of features found here are expected to survive for the isotropic Heisenberg Model. In this last case the excited state gap still vanishes, as in our current example. We also want to emphasize that the general questions of convergence in these methods, which are primarily of a physical and not just a technical mathematical nature, still remain open.

References

- [1] D. Horn and M. Weinstein, Phys. Rev. D **30**, 1256 (1984).
- [2] J. Cioslowski, Phys. Rev. Lett. **58**, 83 (1987).
- [3] J. Cioslowski, Chem. Phys. Lett. **134**, 507 (1987).
- [4] J. Cioslowski, M. Kertesz, P. Surjan, and R. Poirier, Chem. Phys. Lett. **138**, 516 (1987).
- [5] T. Yoshida and K. Iguchi, J. Chem. Phys. **91**, 4249 (1989).
- [6] T. Yoshida and K. Iguchi, Chem. Phys. Lett. **143**, 329 (1988).
- [7] D. Horn, Int. J. Mod. Phys. A **4**, 2147 (1989).
- [8] D. Horn and D. Schreiber, Phys. Rev. D **47**, 2081 (1993).
- [9] D. Horn and E. Klepfish, Bistate t-expansion of the su(2) lattice gauge theory, in *Proceedings of the 25th International Conference on High Energy Physics, vol.1*, edited by K. Phua and Y. Yamaguchi, pages 428–431, Singapore, 1992, World Scientific.
- [10] C. Morningstar, Phys. Rev. D **46**, 824 (1992).
- [11] G. Mathews, N. Snyderman, and S. Bloom, Phys. Rev. D **36**, 2553 (1988).
- [12] D. Horn and G. Lana, Phys. Rev. D **44**, 2864 (1992).
- [13] C. van den Doel and D. Horn, Phys. Rev. D **36**, 3011 (1986).
- [14] W. Massano, S. Bowen, and J. Mancini, Phys. Rev. A **39**, 4301 (1989).
- [15] J. Mancini and W. Massano, Phys. Lett. A **160**, 457 (1991).
- [16] J. Mancini, J. Prie, and W. Massano, Phys. Rev. A **43**, 1777 (1991).
- [17] K. Lee and C. Lo, J. Phys.: Condens. Matter **6**, 7075 (1994).
- [18] K. Lee and C. Lo, Nuovo Cimento **15D**, 1483 (1993).
- [19] C. Lo and Y. Wong, Phys. Lett. A **187**, 269 (1994).
- [20] J. Cioslowski, Int. J. Quantum Chem.: Quant. Chem. Symp. **21**, 563 (1987).
- [21] C. Stubbins, Phys. Rev. D **38**, 1942 (1988).
- [22] P. Knowles, Chem. Phys. Lett. **134**, 512 (1987).
- [23] J. Mancini, Y. Zhou, and P. Meier, Int. J. Quantum Chem. **50**, 101 (1994).
- [24] N. Ullah, Phys. Rev. A **51**, 1808 (1995).

- [25] J. Mancini, W. Massano, J. Prie, and Y. Zhuo, Phys. Lett. A **209**, 107 (1995).
- [26] E. Lieb, T. Schultz, and D. Mattis, Ann. Phys. (NY) **16**, 407 (1961).
- [27] L. Hollenberg, Phys. Lett. A **182**, 238 (1993).
- [28] J. Cioslowski, Phys. Rev. A **36**, 3441 (1987).

Acknowledgements

Part of this work was supported by the Australian Research Council. The author would like to acknowledge encouragement from and valuable discussions with Lloyd C.L. Hollenberg.

List of Figures

1	The error for the ground state energy computed using a D-Páde analysis of $dE(t)/dt$ with a $[M/M + 2]$ approximant, versus M	16
2	The error for the ground state energy computed using an Inversion analysis of $dt(E)/dE$ with $[L/M]$ approximants, versus L and $L + M$ is fixed at various sizes.	17




Figure 1:




Figure 2: